

On a multiple harmonic power series.

Michel Émery *

February 1, 2008

Abstract

The results obtained in the first version are not new.

MSC: 11M35, 33E20, 40A25, 40B05.

Keywords: Multiple harmonic series, Lerch function, Polylogarithm.

New version

The claimed results are not new. I am thankful to Professor Prodinger who kindly pointed out to me references on earlier publication. The lemma of the first version was already published in 1995 as Corollary 3 of [1], and the formulas stated in the abstract of the first version are trivial consequences of it.

References

- [1] K. Dilcher, Some q -identities related to divisor functions, *Discrete Math.* **145** (1995) 83–93.

*IRMA, 7 rue René Descartes, 67084 Strasbourg Cedex, France. emery@math.u-strasbg.fr

On a multiple harmonic power series.

Michel Émery *

February 1, 2008

Abstract

If Li_s denotes the polylogarithm of order s , where s is a natural number, and if z belongs to the unit disk,

$$\text{Li}_s\left(\frac{-z}{1-z}\right) = - \sum_{1 \leq i_1 \leq \dots \leq i_s} \frac{z^{i_s}}{i_1 i_2 \dots i_s}.$$

In particular,

$$\sum_{n \geq 1} \frac{(-1)^{n+1}}{n^s} = \sum_{1 \leq i_1 \leq \dots \leq i_s} \frac{1}{i_1 \dots i_s 2^{i_s}}.$$

MSC: 11M35, 33E20, 40A25, 40B05.

Keywords: Multiple harmonic series, Lerch function, Polylogarithm.

Introduction

Equalities and identities between multiple harmonic series and polylogarithms have been investigated by many authors; see for instance [1] and the references therein. These series usually involve summations over all s -tuples $(i_1, \dots, i_s)$ of natural numbers such that $i_1 < \dots < i_s$, where s is fixed. We shall be concerned with an instance where the summation indices may be equal to each other, that is, a sum over all integers $i_1, \dots, i_s$ verifying $1 \leq i_1 \leq \dots \leq i_s$.

Definition. For $\alpha \in \mathbb{C} \setminus \{-1, -2, \dots\}$ and $s \in \{1, 2, \dots\}$, the Lerch function of order s with shift α is defined by

$$\text{Li}_s^\alpha(w) = \sum_{n \geq 1} \frac{w^n}{(\alpha + n)^s}.$$

The power series converges in the unit disk only, but the analytic function Li_s^α extends to the whole complex plane minus a cut along for instance the

*IRMA, 7 rue René Descartes, 67084 Strasbourg Cedex, France. emery@math.u-strasbg.fr

half-line $[1, \infty)$; see § 1.11 of [2]. When $\alpha = 0$, Li_s^α is just Li_s , the usual polylogarithm of order s . We shall be interested in the values of $\text{Li}_s^\alpha(w)$ in the half-plane $\text{Re}(w) < \frac{1}{2}$ only; they are given by the next proposition (which establishes anew the existence of the analytic extension of Li_s^α to that half-plane).

Proposition. *For $\alpha \in \mathbb{C} \setminus \{-1, -2, \dots\}$ and $s \in \{1, 2, \dots\}$, one has the following power series expansion, which converges for $|z| < 1$:*

$$\begin{aligned} \text{Li}_s^\alpha \left(\frac{-z}{1-z} \right) \\ = - \sum_{1 \leq i_1 \leq \dots \leq i_s} \frac{(i_s - 1)!}{(\alpha + 1)(\alpha + 2) \dots (\alpha + i_s)} \frac{z^{i_s}}{(\alpha + i_1)(\alpha + i_2) \dots (\alpha + i_{s-1})} . \end{aligned}$$

When $\alpha = 0$, the right-hand side becomes much simpler:

Corollary. *Fix $s \in \{1, 2, \dots\}$. For z in the unit disk,*

$$\text{Li}_s \left(\frac{-z}{1-z} \right) = - \sum_{1 \leq i_1 \leq \dots \leq i_s} \frac{z^{i_s}}{i_1 i_2 \dots i_s} .$$

Equivalently, for $\text{Re } w < \frac{1}{2}$,

$$\text{Li}_s(w) = - \sum_{1 \leq i_1 \leq \dots \leq i_s} \frac{1}{i_1 i_2 \dots i_s} \left(\frac{-w}{1-w} \right)^{i_s} .$$

For instance, choosing $z = \frac{1}{2}$ in the corollary gives

$$\sum_{n \geq 1} \frac{(-1)^{n+1}}{n^s} = \sum_{1 \leq i_1 \leq \dots \leq i_s} \frac{1}{i_1 \dots i_s 2^{i_s}} ;$$

as the left-hand side is $(1 - 2^{1-s})\zeta(s)$, and the right-hand one is $\sum a_p / (p2^p)$ with $a_p \leq (1 + \log p)^{s-1}$, this is a reasonably fast series expansion of $\zeta(s)$. More generally, taking $z = \frac{1}{2}$ in the proposition gives a more rapidly convergent series for $\sum_n (-1)^n (\alpha + n)^{-s}$.

Proof of the proposition

Observe first that the right-hand side in the statement of the proposition is a power series $\sum c_p z^p$, with

$$\begin{aligned} |c_p| &\leq \frac{(p-1)!}{|\alpha+1| \dots |\alpha+p|} \sum_{1 \leq i_1 \leq \dots \leq i_{s-1} \leq p} \frac{1}{|\alpha+i_1| \dots |\alpha+i_{s-1}|} \\ &\leq \frac{(p-1)!}{|\alpha+1| \dots |\alpha+p|} \frac{p^{s-1}}{C(\alpha)^{s-1}} , \end{aligned}$$

where $C(\alpha) = \inf \{|\alpha+1|, |\alpha+2|, \dots\} > 0$. By d'Alembert's test, the series $\sum |c_p z^p|$ converges in the unit disk; so does also $\sum c_p z^p$. Consequently, to prove the proposition, it suffices to show that both sides of the claimed identity are equal for $|z|$ small enough. We shall take $|z| < \frac{1}{2}$; for such a z , one has $|-z/(1-z)| < 1$, whence

$$\text{Li}_s^\alpha \left(\frac{-z}{1-z} \right) = \sum_{n \geq 1} \frac{(-1)^n}{(\alpha+n)^s} \left(\frac{z}{1-z} \right)^n = \sum_{n \geq 1} \frac{(-1)^n}{(\alpha+n)^s} \sum_{p \geq n} \binom{p-1}{n-1} z^p.$$

Exchanging the summations gives

$$(*) \quad \text{Li}_s^\alpha \left(\frac{-z}{1-z} \right) = \sum_{p \geq 1} z^p \sum_{n=1}^p \binom{p-1}{n-1} \frac{(-1)^n}{(\alpha+n)^s}.$$

This exchange is licit because, setting $K(\alpha) = 1/\inf_{n \geq 1} |1+\alpha/n| < \infty$, one has the estimate

$$\sum_{p \geq n \geq 1} \left| \binom{p-1}{n-1} \frac{(-1)^n z^p}{(\alpha+n)^s} \right| \leq K(\alpha)^s \sum_{p \geq n \geq 1} \binom{p-1}{n-1} \frac{|z|^p}{n^s} = K(\alpha)^s \sum_{n \geq 1} \frac{1}{n^s} \left(\frac{|z|}{1-|z|} \right)^n,$$

which is finite because $0 \leq |z|/(1-|z|) < 1$.

To establish the proposition, it remains to compute the coefficient of z^p in $(*)$, and more precisely to show that, for all $\alpha \notin \{-1, -2, \dots\}$, $s \in \{1, 2, \dots\}$ and $p \geq 1$,

$$\sum_{n=1}^p \binom{p-1}{n-1} \frac{(-1)^n}{(\alpha+n)^s} = \begin{cases} -\frac{(p-1)!}{(\alpha+1)_p} \sum_{1 \leq i_1 \leq \dots \leq i_{s-1} \leq p} \prod_{r=1}^{s-1} \frac{1}{\alpha+i_r} & \text{if } s \geq 2, \\ -\frac{(p-1)!}{(\alpha+1)_p} & \text{if } s = 1, \end{cases}$$

where $(x)_p$ is the Pochhammer symbol standing for $\prod_{j=0}^{p-1} (x+j)$.

Putting $q = p-1$, $m = n-1$ and $\beta = \alpha+1$, it suffices to prove the following lemma:

Lemma. *For all $\beta \in \mathbb{C} \setminus \{0, -1, \dots\}$ and all integers $q \geq 0$ and $s \geq 1$, one has*

$$\sum_{m=0}^q \binom{q}{m} \frac{(-1)^m}{(\beta+m)^s} = \begin{cases} \frac{q!}{(\beta)_{q+1}} \sum_{0 \leq i_1 \leq \dots \leq i_{s-1} \leq q} \prod_{r=1}^{s-1} \frac{1}{\beta+i_r} & \text{if } s \geq 2; \\ \frac{q!}{(\beta)_{q+1}} & \text{if } s = 1. \end{cases}$$

Proof of the lemma

Call $L(q, \beta)$ and $R(q, \beta)$ the left- and right-hand sides in this statement. First, for $q = 0$ and $q = 1$, the lemma is easily verified:

$$L(0, \beta) = \frac{1}{\beta^s} = R(0, \beta) ;$$

$$L(1, \beta) = \frac{1}{\beta^s} - \frac{1}{(\beta+1)^s} = \frac{1}{\beta(\beta+1)} \sum_{u+v=s-1} \frac{1}{\beta^u} \frac{1}{(\beta+1)^v} = R(1, \beta) .$$

Next, observe that

$$\begin{aligned} L(q+1, \beta) &= \sum_m \binom{q+1}{m} \frac{(-1)^m}{(\beta+m)^s} = \sum_m \left[\binom{q}{m} + \binom{q}{m-1} \right] \frac{(-1)^m}{(\beta+m)^s} \\ &= \sum_m \binom{q}{m} \left[\frac{(-1)^m}{(\beta+m)^s} + \frac{(-1)^{m+1}}{(\beta+m+1)^s} \right] \\ &= L(q, \beta) - L(q, \beta+1) . \end{aligned}$$

The rest of the proof will consist in verifying that, for $q \geq 1$, the right-hand side satisfies the same relation:

$$(**) \quad R(q+1, \beta) = R(q, \beta) - R(q, \beta+1) .$$

When this is done, (**) immediately implies that the property

$$\forall \beta \notin \{0, -1, \dots\} \quad L(q, \beta) = R(q, \beta)$$

extends by induction from $q = 1$ to all $q \geq 2$, thus proving the lemma.

The following notation will simplify the proof of (**): for all integers $n \geq 0$, $t \geq 1$, $a \geq 0$ and $b \geq a$, set

$$f_n = \frac{1}{\beta+n} ; \quad S_a^b(t) = \sum_{a \leq i_1 \leq \dots \leq i_t \leq b} \prod_{r=1}^t f_{i_r} ; \quad S_a^b(0) = 1 ;$$

and remark that, for $t \geq 0$ and $0 \leq a \leq b < c$,

$$S_a^c(t) = \sum_{u+v=t} S_a^b(u) S_{b+1}^c(v) ;$$

similarly, for $1 \leq a \leq b$,

$$S_{a-1}^{b+1}(t) = \sum_{u+v+w=t} S_{a-1}^{a-1}(u) S_a^b(v) S_{b+1}^{b+1}(w) = \sum_{u+v+w=t} f_{a-1}^u S_a^b(v) f_{b+1}^w .$$

Keeping these remarks in mind, the proof of (**) goes as follows. Put $t = s - 1$ and write

$$\begin{aligned} R(q, \beta) - R(q, \beta+1) &= \frac{q!}{(\beta)_{q+1}} S_0^q(t) - \frac{q!}{(\beta+1)_{q+1}} S_1^{q+1}(t) \\ &= \frac{q!}{(\beta+1)_q} \left[\frac{1}{\beta} S_0^q(t) - \frac{1}{\beta+q+1} S_1^{q+1}(t) \right] . \end{aligned}$$

The quantity in square brackets can be rewritten as

$$\begin{aligned}
& f_0 \sum_{v=0}^t S_0^0(t-v) S_1^q(v) - f_{q+1} \sum_{v=0}^t S_1^q(v) S_{q+1}^{q+1}(t-v) \\
&= \sum_{v=0}^t S_1^q(v) (f_0^{t-v+1} - f_{q+1}^{t-v+1}) = \sum_{v=0}^t S_1^q(v) (f_0 - f_{q+1}) \sum_{u+w=t-v} f_0^u f_{q+1}^w \\
&= (f_0 - f_{q+1}) \sum_{u+v+w=t} f_0^u S_1^q(v) f_{q+1}^w = (f_0 - f_{q+1}) S_0^{q+1}(t) \\
&= \frac{q+1}{\beta(\beta+q+1)} S_0^{q+1}(t) .
\end{aligned}$$

Finally, the difference $R(q, \beta) - R(q, \beta+1)$ amounts to

$$\frac{q!}{(\beta+1)_q} \frac{q+1}{\beta(\beta+q+1)} S_0^{q+1}(t) = \frac{(q+1)!}{(\beta)_{q+2}} S_0^{q+1}(t) = R(q+1, \beta) .$$

This proves (**), and at the same time the lemma and the proposition.

Remark. In the particular case when $\alpha = 0$ and $z = \frac{1}{2}$, (*) is exactly formula (4) of J. Sondow [3], obtained by accelerating convergence of the alternating zeta series. What the lemma does is computing the numerator in that formula. To that end, (**) is needed for $\beta \in \{1, 2, \dots\}$ only; but the general proof is just as easy.

References

- [1] D. Bailey, P. Borwein, D.J. Broadhurst, and P. Lisoněk, Special values of multiple polylogarithms, *Trans. Amer. Math. Soc.* **353** (2000) 907–941.
- [2] *Higher Transcendental Functions*, A. Erdélyi editor (McGraw-Hill, New York, 1953).
- [3] J. Sondow, Analytic continuation of Riemann’s zeta function and values at negative integers via Euler’s transformation of series, *Proc. Amer. Math. Soc.* **120** (1994) 421–424.